

# MHC class I expression dependent on bacterial infection and parental factors in whitefish embryos (Salmonidae)

EMILY S. CLARK,\*<sup>1</sup> LAETITIA G. E. WILKINS\*<sup>1</sup> and CLAUS WEDEKIND\*

\*Department of Ecology and Evolution, University of Lausanne, Biophore, 1015 Lausanne, Switzerland

## Abstract

Ecological conditions can influence not only the expression of a phenotype, but also the heritability of a trait. As such, heritable variation for a trait needs to be studied across environments. We have investigated how pathogen challenge affects the expression of MHC genes in embryos of the lake whitefish *Coregonus palaea*. In order to experimentally separate paternal (i.e. genetic) from maternal and environmental effects, and determine whether and how stress affects the heritable variation for MHC expression, embryos were produced in full-factorial *in vitro* fertilizations, reared singly, and exposed at 208 degree days (late-eyed stage) to either one of two strains of *Pseudomonas fluorescens* that differ in their virulence characteristics (one increased mortality, while both delayed hatching time). Gene expression was assessed 48 h postinoculation, and virulence effects of the bacterial infection were monitored until hatching. We found no evidence of MHC class II expression at this stage of development. MHC class I expression was markedly down-regulated in reaction to both pseudomonads. While MHC expression could not be linked to embryo survival, the less the gene was expressed, the earlier the embryos hatched within each treatment group, possibly due to trade-offs between immune function and developmental rate or further factors that affect both hatching timing and MHC expression. We found significant additive genetic variance for MHC class I expression in some treatments. That is, changes in pathogen pressures could induce rapid evolution in MHC class I expression. However, we found no additive genetic variance in reaction norms in our study population.

*Keywords:* *Coregonus*, genetic variation, life history, plasticity, *Pseudomonas fluorescens*, reaction norms

Received 12 August 2012; revision received 16 July 2013; accepted 18 July 2013

## Introduction

The quantity of mRNA transcribed for a particular gene is one of the earliest observable phenotypes (Aubin-Horth & Renn 2009; Hodgins-Davis & Townsend 2009), and like many other phenotypes, can be heavily influenced by environmental conditions (Picard & Schulte 2004; Nath *et al.* 2006; Fisher & Oleksiak 2007; Larsen *et al.* 2007; Croisetiere *et al.* 2010; Debes *et al.* 2012), by genotype (Cavaliere *et al.* 2000; Townsend *et al.* 2003; Debes *et al.* 2012; Runcie *et al.* 2012), and by their interactions (e.g. Landry *et al.* 2006; Li *et al.* 2006; Côté *et al.*

2007; McCairns & Bernatchez 2010; Hodgins-Davis *et al.* 2012). In cases where variation in gene expression has a genetic component, the trait has the potential to quickly evolve in response to changing selection pressures (Schlichting & Pigliucci 1998) if genetic covariance between traits and selection acting on correlated traits do not inhibit the expected evolutionary changes (Merilä *et al.* 2001).

The heritability of a trait is typically not constant but varies across environments (Wilson *et al.* 2006). There appears to be a lack in consistency in the immediate effects that different types of environmental conditions have on the amount of genetic variation that can be observed at a given moment (Hoffmann & Merilä 1999). Unfavourable conditions, that is, situations leading to an immediate reduction in fitness, can either (i)

Correspondence: Claus Wedekind, Fax: +41 21 692 42 65;

E-mail: claus.wedekind@unil.ch

<sup>1</sup>Shared first authorship.

decrease heritable variation, for example, by changing the environmental variance component (Charmanier & Garant 2005) or by preventing an organism from reaching its genetic potential (Merilä & Sheldon 1999); (ii) increase heritable variation (Agrawal *et al.* 2002; Relyea 2005), for example, by amplifying phenotypic differences between genotypes or by lowering the threshold for trait expression and releasing cryptic genetic variation (Gibson & Dworkin 2004; McGuigan & Sgro 2009); or (iii) may have no detectable effect on heritable variation (Pakkasmaa *et al.* 2003; Merilä *et al.* 2004; Clark *et al.* 2013). If the impact of environmental change on trait heritability is likely dependent on both the stressor and trait at hand, and perhaps even varies among types of genes, evaluation on a case-by-case basis seems necessary. For genes of the immune system, whose expression can be strongly dependent on the biotic environment (Frost 1999), understanding the impact of ecological stressors on heritable variation for trait means and the norms of reaction (i.e. the function that relates the phenotypes that can be produced by one genotype across environments; Pigliucci 2001) is important, as it would indicate a population's ability to evolve in response to parasites and infectious diseases. Parasites and pathogens are ubiquitous (Windsor 1998), but pressures from microbial organisms are often expected to increase in wild populations, due to pollution, habitat degradation and climate change (Harvell *et al.* 1999; Daszak 2000; Dobson & Foufopoulos 2001).

We have studied the effects of pathogen treatment on the expression of a major histocompatibility complex (MHC) gene, that is, a key component of the adaptive immune system of vertebrates. As a vertebrate model, we chose a wild population of the lake whitefish *Coregonus palaea* (Salmonidae) (Kottelat & Freyhof 2007). *Coregonus palaea* is an iteroparous, fast-growing, lake-dwelling Alpine whitefish that feeds mainly on zooplankton and insect larvae, and spawns once a year during a few days in early winter. Average body length of mature fish at the spawning place is 383 mm (SD = 27) as determined from a random sample of 30 fish. Like all Alpine whitefish, *C. palaea* has external fertilization and shows no parental care. Fertilized eggs simply cascade onto the lake floor where embryo development takes place over a period of approximately 300 degree days. Embryos are therefore, over several weeks, exposed to microbes and further environmental stressors. Whitefish are excellent models for ecological and quantitative genetic studies because individuals produce large amounts of gametes that can be collected for experimental *in vitro* fertilization, including large-scale full-factorial breeding designs. Embryos can be reared singly or in groups under very controlled conditions and monitored until hatching (Wedekind *et al.* 2001,

2008; Wedekind & Müller 2005; Clark & Wedekind 2011). Salmonid embryos usually show high survivorship under benign laboratory conditions (e.g. von Siebenthal *et al.* 2009; Clark *et al.* 2013), but mortality rates in the wild can be high (Stelkens *et al.* 2012). While a number of factors including pollution (Heintz *et al.* 1999), predation (Phillips & Claire 2011), temperature (Tang *et al.* 1987) and oxygen deprivation (Silver *et al.* 2011) can contribute to this mortality, pathogens likely exert strong selection pressures (Schreck *et al.* 2001; Arkush *et al.* 2002), and seemingly benign microbial symbiotic communities can quickly turn virulent under altered environmental conditions (Jacob *et al.* 2010; Wedekind *et al.* 2010). The ecological relevance of microbial pathogens is corroborated by the observations that salmonid embryos have evolved early immunological defence mechanisms (see below). They are also able to perceive waterborne cues from microbial infections and to switch life history strategies accordingly (Wedekind 2002; Pompini *et al.* 2013).

We chose to use two isolates of the opportunistic fish pathogen *Pseudomonas fluorescens* (Austin & Ausin 1999) as microbial stressors. This bacterium is found widely in the aquatic environment (Austin & Ausin 1999; Spiers *et al.* 2000) and has been associated with disease pathologies not only in adult fish (Zhang *et al.* 2009), but also in embryos of whitefish (Wedekind *et al.* 2001; von Siebenthal *et al.* 2009) and brown trout (Clark *et al.* 2013; Pompini *et al.* 2013). Moreover, previous studies suggest that this bacterium has strain-dependent virulence effects on salmonid embryos, with certain isolates directly increasing embryonic mortality (von Siebenthal *et al.* 2009; Pompini *et al.* 2013), and other isolates causing sublethal effects, that is, delayed hatching and reduced growth (Clark *et al.* 2013). We assessed whether challenge with both 'high' (i.e. induced mortality) and 'low' (i.e. delayed hatching) virulence strains resulted in similar MHC expression patterns.

The extent to which embryos can mount an immune response against pathogens at this developmental stage is not clear yet. The immune system of teleost fish is generally thought to only become completely functional after hatching (Fischer *et al.* 2005; Zapata *et al.* 2006). However, the timing of maturation may vary between species (Magnadottir 2006; Mulero *et al.* 2007). Fischer *et al.* (2005) found that MHC class I transcription begins shortly after fertilization in rainbow trout (*Oncorhynchus mykiss*). Mortality of salmonid embryos has been demonstrated to be both MHC-allele-specific (Pitcher & Neff 2006) and dependent on nucleotide diversity at the MHC loci (Evans *et al.* 2010a), and mortality during an epidemic of a nonspecified strain of *P. fluorescens* has led to a significant shift of MHC-allele frequencies within one of seven families of another lake whitefish

(Wedekind *et al.* 2004). Hence, the MHC can already play a role in determining the susceptibility of salmonids to pathogens at late embryogenesis, either through direct pathogen-binding action or via pleiotropic interactions and/or linkages with other viability genes. Notably, both classes of the MHC are not necessarily ideal candidates for an expression study during embryonic development. In teleost fish, class I and class II genes are in separate linkage groups (Sato *et al.* 2000), and the beginning of transcription is not always synchronous (Rodrigues *et al.* 1998). We, therefore, first examined whether transcripts of both MHC classes were detectable in the whitefish embryos in an effort to identify a suitable locus for our study. We then assessed whether bacterial infection changed MHC expression patterns. As we employed a full-factorial breeding design, we were able to provide first estimates of the heritable variation for gene expression and examine whether this varied according to environmental conditions. Finally, we determined whether there was heritable variation for gene expression reaction norms and assessed whether there was a relationship between MHC gene expression and embryo mortality or time to hatching.

## Materials and methods

### *Whitefish sampling and rearing of embryos*

Large-type adult whitefish were caught from their spawning grounds in Lake Geneva with gill nets and stripped of their gametes. These gametes were subsequently used for full-factorial *in vitro* fertilizations following the methods described in von Siebenthal *et al.* (2009). Embryos were distributed singly into 24-well plates (Falcon, Becton Dickinson) in a blockwise design and from then on stored in a 6.5 °C climate chamber. Without additional challenge, this method for rearing embryos typically results in survival rates close to 100% (e.g. von Siebenthal *et al.* 2009; Clark *et al.* 2013; Pompini *et al.* 2013). Embryos were monitored weekly with a light table (Hama professional, LP 555) and a stereo zoom microscope (Olympus SZX9) until the start of hatching, at which point they were monitored daily. Thirteen offspring of 16 different sibships (resulting from a 4 × 4 cross) for each of the four treatments were randomly selected for subsequent work ( $N_{\text{total}} = 13 \times 16 \times 4$  singly reared embryos).

### *Identification of pseudomonad sequence differences*

The two *Pseudomonas fluorescens* isolates used were the 'high virulence strain' DSM 50090 ('PF1') that had been linked to mortality in whitefish embryos (von

Siebenthal *et al.* 2009) and a 'low virulence strain' ('PF2') that had been isolated from whitefish gills and had been observed to cause little embryonic mortality, but to delay hatching of smaller larvae in both brown trout (Clark *et al.* 2013) and whitefish (E. Clark, unpublished data). PF2 had been collected by swabbing gills with Amies agar gel transport swabs, followed by elution of bacteria into phosphate-buffered saline. A  $10^{-2}$  dilution had been plated onto King's B agar to facilitate isolation of fluorescent pseudomonads. After incubation for 48 h, a colony was randomly selected and restreaked three times to obtain a pure culture. To compare sequences of the two pseudomonads (and to confirm successful identification), DNA was first isolated from both using the GenElute™ Bacterial Genomic DNA Kit, according to the manufacturer's instructions (Sigma-Aldrich). PCR was performed with a *P. fluorescens*-specific primer set, 16SPSEfluF and 16SPSER (Scarpellini *et al.* 2004), which amplifies a 850-bp fragment of the 16S rRNA. The PCR was performed in a total volume of 25 µL and contained 50 ng bacterial genomic DNA, 2.5 µL of 10× PCR buffer, 400 µM of each dNTP, 2.5 mM of MgCl<sub>2</sub>, 0.6 µM of each primer and 0.625 U of Taq polymerase (Invitrogen, Carlsbad, CA). The thermal profile was modified from Scarpellini *et al.* (2004): 3 min at 94 °C; 35 cycles of 94 °C for 30 s, 50 °C for 30 s, 72 °C for 1 min and a final extension at 72 °C for 10 min. Following the PCR, the amplified products were purified with the Wizard® SV Gel and PCR Clean-Up System (Invitrogen) and sequenced in the forward and reverse directions with 16SPSEfluF/16SPSER on a ABI Prism 3100 genetic analyzer (Applied Biosystems). Sequences were edited and aligned with Geneious Pro™, version 5.3.4 (Biomatters). Alignment of PF1 and PF2 showed a base pair difference (A-G) at position 733, relative to a reference strain (strain CCM 2115; GenBank: DQ207731.2). To confirm the one base pair difference in the 16S rRNA fragment between the two pseudomonads, two further PCRs were performed as described above, and the resulting fragments were sequenced.

### *Preparation of bacterial inocula and treatment of embryos*

Once embryos had reached the late-eyed stage (208 degree days), two flasks, each containing 100 mL of nutrient broth (3 g meat extract, 5 g bactopectone, 1 L distilled H<sub>2</sub>O), were inoculated with either PF1 or PF2. As the two strains were observed to have different optimal growth temperatures, they were incubated at 30 °C or 22 °C, respectively, for 36 h on shakers until reaching the exponential growth phase. The bacteria were transferred to 50-mL conicals and spun at 4000 rpm for

15 min. The resulting pellet was washed three times and resuspended in sterile water, standardized according to OECD guidelines (OECD 1992). A Helber counting chamber was used to assess bacterial concentrations (see: Bast 2001, p. 280–285). The suspension was then diluted such that inoculation with 100  $\mu\text{L}$  would achieve a concentration of  $10^8$  bacterial cells/mL in the wells. Prior to inoculation, nutrient broth was added to the suspension to encourage bacterial growth, resulting in a 1:1000 concentration in the wells. Thirteen replicates of every parental combination received PF1 and 13 received PF2. The remaining plates served as controls and were either sham-treated with sterile standardized water ( $N = 13$ ) or were inoculated with nutrient broth ( $N = 13$ ; 1:1000 dilution per well).

#### *RNA preservation and extraction from embryos*

Three embryos per sibship were sampled from each treatment group at 48 h postinoculation, and the remaining ten embryos per sibship and treatment group were monitored for survival and time until hatching. All samples were placed in RNAlater<sup>®</sup> (Ambion, Austin, TX), stored overnight at 6.5 °C and then at –20 °C for long-term storage. Embryos were individually homogenized with a mixer mill (MM300; Retsch, Düsseldorf, Germany) using six tungsten beads (3 mm), five silica beads (1.5 mm) and 0.4 g silica powder (0.2 mm) (Qiagen, Valencia, CA). Total RNA was extracted using the RNeasy Mini Kit (Qiagen) according to the manufacturer's instructions. RNA was eluted with 40  $\mu\text{L}$  of RNase-free water, and an additional DNase treatment was performed in which the following was added to each sample: 40 u/ $\mu\text{L}$  RNasin<sup>®</sup> (Promega, Madison, WI, USA), Tris-HCl (pH 7.5, 1 M),  $\text{MgCl}_2$  (100 mM), KCl (2.5 M), DTT (100 mM), and DNase I (10 u/ $\mu\text{L}$ ) (Roche, Mannheim, Germany). Each sample was incubated for 15 min at 37 °C, followed by a phenol–chloroform extraction and ethanol precipitation. The resulting pellet was resuspended in 10  $\mu\text{L}$  of RNase-free water. RNA integrity was verified by measuring absorbance at 260–280 nm, and a random set of 12 extractions was analysed on an Agilent 2100 Bioanalyzer (Agilent Technologies, Waldbronn, Germany) to confirm that the ratio of 28s/18s rRNA was close to two.

#### *Reverse transcription, PCR of cDNA and real-time quantitative PCR*

Total extracted RNA was reverse-transcribed using the SuperScript<sup>™</sup> III First-Strand Synthesis System (Invitrogen) and random hexamers in a 25  $\mu\text{L}$  reaction according to the manufacturer's protocol. To assess whether MHC class I and class II transcripts were detectable in

the whitefish embryos, a PCR was conducted to amplify the two different genes from cDNA of whole embryos ( $N = 16$ ) and, as a positive control, from spleens of adult whitefish ( $N = 8$ ; RNA preserved and extracted as described above). The PCR was performed in a total volume of 25  $\mu\text{L}$  and contained 10–100 ng of cDNA, 2.5  $\mu\text{L}$  of 10 $\times$  PCR buffer, 400  $\mu\text{M}$  of each dNTP, 2.5 mM of  $\text{MgCl}_2$ , 0.6  $\mu\text{M}$  of each primer (Table 1) and 0.625 U of Taq polymerase (Invitrogen). As a negative control, water was added instead of cDNA. As an additional control, the two genes were verified to successfully amplify from genomic DNA of whole embryos ( $N = 8$ ) and also from fin clips of adult whitefish ( $N = 8$ ). DNA had been extracted using the DNeasy Blood and Tissue Kit (Qiagen) (according to the manufacturer's instructions). The thermal profile for both cDNA and DNA PCR consisted of: 3 min at 94 °C; 32 cycles of 94 °C for 30 s, 56 °C for 30 s, 72 °C for 30 s and a final extension at 72 °C for 10 min. Amplified products were subsequently run on a 2% agarose gel, stained with ethidium bromide and visualized under UV light.

Expression of MHC class I, along with four housekeeping genes (G6PD, GAPDH,  $\beta$ -actin and NADH), was analysed with real-time qPCR using primers developed for salmonids (Table 1). Primer efficiencies were verified for each pair with fourfold serial dilutions of cDNA. Each 10  $\mu\text{L}$  reaction contained: 2  $\times$  SYBR<sup>®</sup> Green PCR Mastermix (Roche Diagnostics, Mannheim, Germany), 0.3  $\mu\text{M}$  of each primer and 1  $\mu\text{L}$  of cDNA. Three technical triplicates of each reaction were run on a 7900HT Fast Real-Time PCR Sequence Detection System (Applied Biosystems) with the following conditions: 94 °C for 3 min, followed by 40 cycles of 94 °C for 10 s, 60 °C for 25 s and 72 °C for 30 s. No enzyme controls (NEC) and no template controls (NTC) from the cDNA reaction were run concurrently to test for genomic DNA contamination, and melting curve analysis was used to confirm primer specificity.

The stability of the four candidate reference genes was examined with the method by Vandesompele *et al.* (2002), using the 'ReadqPCR' and the 'NormqPCR' packages (Perkins *et al.* 2012). All four candidate reference genes had been shown to be reliable housekeeping genes in previous expression studies in salmonids (Brzuzan *et al.* 2005, 2007, 2009; Olsvik *et al.* 2005; Benedetto *et al.* 2011). Expression of G6PD and GAPDH, but not  $\beta$ -actin or NADH, was confirmed to remain constant across our treatments, therefore meeting the stability criteria for reference genes. All samples were subsequently normalized to the geometric mean of G6PD and GAPDH, and relative expression of mRNA was determined using the  $2^{-\Delta\Delta\text{CT}}$  method (Livak & Schmittgen 2001; Schmittgen & Livak 2008).

**Table 1** Primers used in the present study

Gene	Forward primer 5'-3'	Reverse primer 5'-3'
GAPDH*	ATG ACC ACT CCA TCT CCG TAT TC	ACG ACG TAA TCG GCA CCG
G6PD†	CCC TAT ATG AAG GTG GCA GAC TCT	GGC GTA CTT CCC ACT GAC ATA AG
β-actin‡	GTG GCG CTG GAC TTT GAG CA	ACC GAG GAA GGA GGG CTG GA
NADH*	CAT CAC CAT CGC ACT ATC CA	CCT CCT TGG GTT CAC TCG TA
MHC class I§	TGT GGC TGT GGG GAT GGT GGA	TTT GGG CAC CGC TCT CTG GC
MHC class II¶	ATG TTT TCC TTT TAG ATG GAT ATT TT	AGC CCT GCT CAC CTG TCT TA

\*Atlantic salmon (Olsvik *et al.* 2005).

†Rainbow trout (Benedetto *et al.* 2011).

‡Whitefish (Brzuzan *et al.* 2007).

§Whitefish, modified from Binz *et al.* (2001).

¶Salmonids (Pavey *et al.* 2011).

### Cloning and sequencing of parental genotypes

The gene of interest from each parent was cloned (Table 2) to confirm amplification of the desired products in the qPCRs and the presence of two coamplified loci in the MHC class I sequences (Binz *et al.* 2001). Cloning was performed with the TOPO TA Cloning® Kit (Invitrogen) using the pCR®2.1-TOPO vector and One Shot®TOP10 chemically competent cells. Between seven and 14 positive clones per individual were selected and amplified with M13 forward (5'-GTA AAA CGA CGA CCA G-3') and reverse (5'-CAG GAA ACA GCT ATG AC-3') primers with the following amplification profile: 94 °C for 2 min, followed by 33 cycles of 94 °C for 15 s, 54 °C for 15 s, and 72 °C 15 s, and a final 10-min extension at 72 °C. Reactions were performed in a total volume of 15 µL and contained: 10 µM of each primer, 1.5 µL of 10× PCR buffer, 0.5 µM of each dNTP, 0.08 µL Taq™ DNA polymerase, 10 ng of cloned DNA and water. Blank PCRs were included as controls. Amplified products were purified with the Wizard® SV Gel and PCR Clean-Up System (Invitrogen) and sequenced in the forward direction with M13 on an ABI Prism 3100 genetic analyzer (Applied Biosystems). Each allele was confirmed to be present in at least three clones to avoid cloning artefacts (Lenz & Becker 2008).

### Statistical analysis

Mortality was analysed as a binomial response variable in general linear mixed effect models (GLMM), while hatching time and MHC class I expression were analysed as continuous response variables in linear mixed effect models (LMM). Prior to analysis, gene expression values, normalized to the housekeeping genes (i.e. ΔCt), were converted to the linear form with the 2<sup>-ΔCt</sup> calculation (Livak & Schmittgen 2001; Schmittgen & Livak 2008) and were log-transformed to avoid deviations from normality. Treatment was entered as a fixed effect,

**Table 2** Observed MHC I genotypes. Primers amplified a 68-bp fragment of the MHC gene

Individual	Alleles	GenBank accession number
Male 1	Cosp-A1-H-6/8*	AF213305/AF213308
	Cosp-A1-H-1	AF213306
Male 2	Cosp-A1-H-15	AF213293
Male 3	Cosp-A1-H-6/8*	
	Cosp-A1-H-1	
	Cosp-A1-H-10	AF213296
Male 4	Cosp-A1-H-6/8*	
	Cosp-A1-H-1	
Female 1	Cosp-A1-H-6/8*	
	Cosp-A1-H-1	
	Cosp-A1-H-10	
Female 2	Cosp-A1-H-11	AF213303
	Cosp-A1-H-1	
Female 3	Cosp-A1-H-15	
	Cosp-A1-H-10	
Female 4	Cosp-A1-H-6/8*	
	Cosp-A1-H-15	
Female 4	Cosp-A1-H-1	
	Cosp-A1-H-11	

\*Matches to either Cosp-A1-H-6 or Cosp-A1-H-8 (Binz *et al.* 2001).

while sire and dam were entered as random effects. While dam effects encompass both genetic and maternal environmental effects, sire effects represent one-quarter of the additive genetic variance, assuming that epistatic effects are negligible (Lynch & Walsh 1998). Due to the fact that we had low replicate numbers (i.e. three) per full-sib family for the expression analysis (see Table S1, Supporting Information for mean expression levels per full-sib family and their associated variances), we did not include the sire × dam interaction as a random effect in the models. In the case of mortality and hatching time, the interaction term was also not included in the reference models, as it was not found to improve

model fit to the data in any treatment. For the analysis, the nutrient broth treatment was used as the baseline control, as both bacterial treatments received the same concentration of supplemental nutrients. While the addition of nutrients did not significantly elevate mortality in comparison with untreated embryos (1.8% increase; GLMM:  $Z = 0.96$ ,  $P = 0.34$ ), it did result in embryos hatching, on average, 1 day later (LMM:  $T = 2.04$ ,  $P = 0.04$ ).

To assess the importance of each effect, a reference model incorporating all relevant terms was compared with a model lacking the effect of interest. To examine the importance of interaction terms, a model incorporating the interaction was compared with the reference model. Akaike's information criteria (AIC), which provide a measure of model fit and model complexity (lower values indicate a better fit to the data) and likelihood ratio tests (LRT) were used to compare model fits. For models examining the importance of interaction terms (i.e. treatment  $\times$  dam and treatment  $\times$  sire), two models were run, one comparing the control with PF1, and one comparing it with PF2. As the two isolates differed in their virulence characteristics, we assessed for each pseudomonad separately whether a random slope-intercept term improved model fit.

To test for an effect of gene expression and treatment  $\times$  gene expression on time until hatching, an analysis of covariance (ANCOVA) was performed. Average hatching time per full-sib family (by treatment) was entered as the dependent variable, with treatment and mean expression per full-sib family entered as independent variables. To examine the effect of gene expression on embryonic mortality, a general linear model (GLM) with a binomial distribution was used, with the proportion of dead embryos (of the 10 initial replicates) per full-sib family entered as the response variable, and mean expression per full-sib family entered again as the independent variable. For this GLM on embryo mortality, we only looked within PF1, as survival in the control and in PF2 was close to 100%.

All analyses were done in the R environment (R Development Core Team 2011), using mainly the lme4 package (Bates *et al.* 2011). The MCMCglmm package (Hadfield 2010) was used as a means of verifying the significance of the treatments and the random effects in explaining variation in offspring phenotypes, as well as obtaining highest posterior density confidence intervals for the treatments and the random terms (see Supplementary Materials for details).

## Results

Transcripts of MHC class I were detected in all whitefish embryos ( $N = 16$ ) and all adults ( $N = 8$ ), while

MHC class II transcripts were only found in the tissue samples of the 8 adults (see Fig. S1, Supporting Information for amplification on agarose gels). Treatment with both *Pseudomonas fluorescens* isolates resulted in decreased MHC class I gene expression (Fig. 1; Table 3; Table S2, Supporting Information). While mortality was only significantly higher in PF1-treated embryos, hatching time was significantly delayed in both PF1 and PF2 (Fig. 2; Table 3; Table S2, Supporting Information). Hatching time in PF1 was still significantly delayed after controlling for the likely confounding effects of potential nonrandom mortality in this treatment (Fig. S2, Supporting Information).

We found significant additive genetic variance for MHC class I expression in the control and the PF1-treated groups, but not in the PF2-treated embryos (Table 4A; see Table S3, Supporting Information for the alternative MCMCglmm model). Analogous patterns could be observed with respect to the dam effects (Table 4A). Significant sire and dam effects on embryonic mortality were found after treatment with the virulent pseudomonad (PF1), but not in the PF2-treated embryos or the control group (Table 4B, Table S3, Supporting Information). In contrast, we observed significant sire effects on hatching time in the control and PF2, but not in PF1-treated embryos (Table 4C, Table S3, Supporting Information). Dam effects on hatching time was significant in PF2, but in neither of the other two treatments.

While we found evidence of additive genetic variance for MHC class I expression, embryonic mortality and hatching time in certain environments, we found no indications of heritable variance for their reaction norms, as indicated by the lack of sire  $\times$  treatment interaction effects (Fig. 3; Table 3, see Table S4, Supporting Information for the alternative MCMCglmm model). In contrast, we did find evidence that the virulence of a given pseudomonad was modulated by dam effects, as





Fig. 1 Mean MHC class I gene expression per treatment ( $\pm$ SE).

**Table 3** Likelihood ratio tests on mixed model logistic regressions on MHC class I expression, embryonic mortality and hatching time. Treatment was entered as a fixed effect [two levels including the control and PF1 (A) or the control and PF2 (B)], while sire and dam were entered as random effects. To test the effect of treatment, a reduced model was compared with the reference model (in bold). To test for interaction effects, a model incorporating the term was compared with the reference model. Akaike's information criteria (AIC), which provide a measure of model fit and model complexity (lower values indicate a better fit to the data), and likelihood ratio tests (LRT) were used to compare model fits ( $\chi^2$ )

Model	Effect tested	MHC I expression				Embryonic mortality				Hatching time			
		AIC	$\chi^2$	DF	P	AIC	$\chi^2$	DF	P	AIC	$\chi^2$	DF	P
<b>(A) Control vs. PF1</b>													
<b>t + s + d</b>		66.3		5		135.1		4		1757.2		5	
s + d	t	70.5	6.2	4	0.01	160.0	26.8	3	<0.001	1785.5	30.3	4	<0.001
t + t × s + d	t × s	69.7	0.6	7	0.73	139.1	0.1	6	0.98	1761.2	0	7	1
t + s + t × d	t × d	69.5	0.8	7	0.68	123.8	15.3	6	<0.001	1760.1	1	7	0.59
<b>(B) Control vs. PF2</b>													
<b>t + s + d</b>		40.4		5		82.2		4		1828.9		5	
s + d	t	42.5	4.1	4	0.04	80.7	0.5	3	0.48	1948.0	121.1	4	<0.001
t + t × s + d	t × s	42.4	2	7	0.36	86.2	0	6	1	1831.6	1.3	7	0.52
t + s + t × d	t × d	42.6	1.8	7	0.4	85.8	0.3	6	0.84	1820.8	12.1	7	0.002

t, treatment; s, sire; d, dam; t × s, treatment × sire; t × d, treatment × dam interactions; DF, degrees of freedom.



**Fig. 2** Treatment effects on (A) embryo mortality and (B) time until hatching (means  $\pm$  SE).

suggested by the significant dam  $\times$  treatment effects on embryo mortality in response to PF1 (Table 3A; Table S4, Supporting Information) and hatching time in response to PF2 (Table 3B; Table S4, Supporting Information).

We found no relationship between mean MHC gene expression and survival per full-sibship in PF1, that is, in the one bacterial strain that increased embryo mortality (GLM:  $T = 0.72$ ,  $P = 0.47$ ). However, there was an overall effect of gene expression on hatching time (ANCOVA:  $F_{1,41} = 5.0$ ,  $P = 0.03$ ), with faster hatching embryos expressing less MHC (Fig. 4). No interaction between treatment and gene expression was found (ANCOVA:  $F_{2,41} = 0.90$ ,  $P = 0.41$ ): that is, the link between MHC expression and hatching time was similar in all treatment groups.

## Discussion

We have described the experimental infection of whitefish embryos with two isolates of the opportunistic

pathogen, *Pseudomonas fluorescens*, and resultant effects on MHC expression, embryo mortality and hatching time. A full-factorial experimental design was used to disentangle treatment from sire and dam effects on embryo traits, to assess whether and how environmental stress affects heritable variation for these traits and to test for parental effects on embryo reaction norms. In addition, comparisons of full-sib families allowed testing whether MHC expression was linked to embryo survival or timing of hatching.

### MHC expression in whitefish embryos and the effects of pathogen challenge

Transcripts of MHC class I, but not II, were detectable in the whitefish embryos under our experimental conditions. While our observations do not entirely exclude the possibility that MHC class II is expressed in the embryos at this point in development, as biases can be introduced during reverse transcription (Bustin & Nolan 2004), they nevertheless suggest that class I

expression perhaps begins before class II. A study by Evans *et al.* (2010a) similarly suggested that the MHC class I pathway plays a significant role in survival before the MHC class II pathway, as their results provide strong support for a nucleotide diversity advantage at the MHC class I during the embryonic stage in another salmonid, the Chinook salmon (*O. tshawytscha*). Notably, Wedekind *et al.* (2004) found the allelic specificity on an MHC class II locus to influence survival until hatching in another whitefish during an epidemic with *P. fluorescens*. Pitcher & Neff (2006) also found evidence of MHC class IIB allele- and genotype-dependent survivorship during early developmental stages (i.e. through endogenous feeding) in the Chinook salmon. These findings, combined with the

present ones, suggest that genes of MHC class I and class II can both be expressed prior to hatching, but that during the course of embryogenesis, class I genes are actively transcribed before class II.

Traditionally, MHC class I has been associated with the presentation of endogenous antigens (i.e. viruses and obligate intracellular bacteria), with class II binding proteins of exogenous origins. However, it is now well established that a significant amount of cross-over occurs between the two pathways (Kovacsovic-Bankowski & Rock 1995; Norbury *et al.* 1995; Yewdell *et al.* 1999; Ackerman & Cresswell 2004). Moreover, recent studies in both Atlantic cod (*Gadus morhua*) (Star *et al.* 2011) and pipefish (*Syngnathus typhle*) (Haase *et al.* 2013) have demonstrated a complete absence of MHC class II, suggesting that MHC class I can play an integral role in teleost adaptive immunity.


We found expression of MHC class I in whitefish embryos to vary according to environmental conditions. Specifically, treatment with the two different pseudomonads resulted in an average decrease in expression 48 h postinoculation. A very similar expression phenotype was elicited, despite the fact that the two isolates showed different virulence characteristics (i.e. one induced mortality, as in von Siebenthal *et al.* 2009; while the other only delayed hatching, as in Clark *et al.* 2013). It is possible that a different pattern may have emerged had we sampled embryos at other time points postinoculation, because gene expression can be very dynamic. However, the observed down-regulation of MHC expression confirms a number of previous observations in other fish: Koppang *et al.* (1999) found down-regulation of MHC class I following immunostimulation with lipopolysaccharide in Atlantic salmon (*Salmo salar*), and Reyes-Becerril *et al.* (2011) reported a reduction in MHC class II expression in gilthead seabream (*Sparus aurata*) after infection with *Aeromonas hydrophila*. While these studies did not specifically investigate the reasons for this down-regulation, one explanation could involve pathogen-mediated suppression of transcription.

**Table 4** REML estimates of variance components ( $V_{\text{Sire}}$ ,  $V_{\text{Dam}}$  and  $V_{\text{Res}}$ ) for MHC class I expression, embryonic mortality and hatching time in each treatment (control, PF1 and PF2). Numbers in parentheses indicate percent of total variance explained by each separate component. The significance of each variance component was determined by comparing a mixed effect model incorporating all effects of interest to one lacking it (see Materials and methods)

	$V_{\text{Sire}}$	$V_{\text{Dam}}$	$V_{\text{Res}}$
(A) MHC I expression			
Control	0.06 (40.0)***	0.02 (13.3)*	0.07 (46.7)
PF1	0.05 (27.7)**	0.03 (16.7)*	0.10 (55.6)
PF2	0.02 (22.2)	0.01 (11.1)	0.06 (66.7)
(B) Embryonic mortality			
Control	0 (0)	0 (0)	1 <sup>†</sup>
PF1	2.1 (21.6)**	6.6 (68.0)***	1 <sup>†</sup>
PF2	0 (0)	0.5 (33.3)	1 <sup>†</sup>
(C) Hatching time			
Control	3.1 (18.2)***	0.4 (0.02)	13.5 (79.4)
PF1	2.6 (6.4)	0.8 (1.9)	37.1 (91.7)
PF2	5.8 (17.0)***	3.6 (10.5)**	24.8 (72.5)

\* $P < 0.05$ ; \*\* $P < 0.01$ , \*\*\* $P < 0.001$ .

<sup>†</sup>Because mortality was a binomial response variable, residual variance was set to one.



**Fig. 3** Reaction norms of MHC class I gene expression across (A) control and exposure to PF1 and (B) control and exposure to PF2. Lines correspond to means per paternal sibgroup ( $N = 4$ ).






Fig. 4 Days to hatching vs. MHC class I expression. Points represent means across treatments per full-sib family ( $N = 16$ ).

Specifically, a number of bacteria and viruses seem capable of inhibiting expression of MHC as a means of immune evasion (e.g. Finlay & McFadden 2006; Antoniou & Powis 2008; Lapaque *et al.* 2009).

Another possible explanation for the observed down-regulation of MHC in our experiment is that a trade-off exists with other metabolic functions or immune pathways (Lochmiller & Deerenberg 2000). Life history theory predicts that the immune response is a trait whose expression exacts an important cost on the organism (Moret & Schmid-Hempel 2000). Consequently, resource investment in this response will come at the expense of other traits (e.g. growth or reproduction) (Norris & Evans 2000; Schmid-Hempel 2003). In the context of our experiments, we found an overall positive relationship between MHC gene expression and hatching date. While fish, among other vertebrates, can alter hatching age to mitigate the fitness consequences of environmental stressors (e.g. pathogens and predators) (Warkentin 2011), late hatching is typically selected against in salmonids (Koho *et al.* 1991; Einum & Fleming 2000; Skoglund *et al.* 2012) and has been shown to be associated with reduced larval survival in another coregonid (*C. albula*) (Koho *et al.* 1991). Decreased transcription of MHC could, therefore, potentially reflect a strategic decision to invest resources into growth and an earlier life history transition. Somewhat paradoxically, embryos raised in benign experimental conditions still managed to hatch earlier and express the most MHC. However, as they were not subjected to the burden of a pathogen challenge, they perhaps were in a better position to attain optimal phenotypes with respect to both traits. A comprehensive examination of the transcriptional changes of a wider array of genes, involved in both immunity and development, could help clarify the mechanisms behind this possible trade-off. Such an examination could also help shed light on whether MHC class I expression and

developmental time are both affected by further factors that explain the somewhat counterintuitive observation that MHC class I expression declines with infection, hatching is delayed by infection, but there is a positive correlation between MHC class I expression and hatching date within each treatment group.

#### *Components of phenotypic variation and the consequences of environmental stress*

We found significant heritable variation for MHC class I expression and hatching time under benign experimental conditions. As both of these traits are closely tied to fitness, one would traditionally expect them to be characterized by reduced additive genetic variance due to directional selection (Mousseau & Roff 1987). However, a number of studies have demonstrated that significant heritable variation is often maintained in fitness-related traits (e.g. Laurila *et al.* 2002; Jacob *et al.* 2007, 2010), with high residual variance sometimes giving the impression of depleted heritable variation (Houle 1992). On the contrary, little additive genetic variance was observed for embryonic mortality under these conditions, although our power to detect such variation was limited by the overall lack of mortality.

We found that the significance of the sire effect changed for some of the monitored traits and not for others in the *Pseudomonas* treatments, suggesting environmental dependencies. Heritable variation for gene expression and for hatching time seemed to decrease after inoculation with one of the *Pseudomonas* isolates; however, both isolates did not seem to have the same effect on heritable variation for a given trait, suggesting that different strains of the same bacterium can differentially affect a trait's evolutionary potential. Somewhat analogous declines in the heritability of expression have been observed in another set of immune genes, that is, cytokines, postimmunostimulation with a *Vibrio* vaccine in Chinook salmon (Aykanat *et al.* 2012). Heritable variation for embryonic mortality seemed to increase under stressful conditions (i.e. in PF1), potentially due to a release of cryptic genetic variation that, under different circumstances, would be phenotypically neutral (Gibson & Dworkin 2004; McGuigan & Sgro 2009). These observations support the view that the effect of environmental stressors on additive genetic variance is not only trait dependent, but also dependent on the stressor at hand (for other examples, see Hoffmann & Merilä 1999; Laugen *et al.* 2005). However, none of the possible environmental dependencies on heritability could be confirmed in pairwise comparisons between the controls and the pathogen strain (i.e. the sire  $\times$  treatment interaction terms were never statistically significant).

As with sire effects, the importance of dam effects on offspring phenotype can also vary according to the ecological conditions (Einum & Fleming 1999; Laugen *et al.* 2005 and references therein). In the case of MHC class I expression, the significance of dam effects appeared to remain stable after treatment with one isolate of *P. fluorescens* (PF1), but to decrease following inoculation with the second (PF2). No dam  $\times$  treatment interaction on gene expression was found, suggesting that the decrease in transcription across environments was uniform across females. The significance of dam effects on embryonic mortality and hatching time increased under certain pathogen conditions, as confirmed also in significant dam  $\times$  treatment interaction terms in the pairwise comparisons between the controls and the pathogen strains. Maternal sibgroups must have varied in their response to treatment either due to characteristics of the maternal environmental contributions [e.g. immune compounds (Magnadottir 2006)] or due to genetic effects.

#### *Heritable variation for embryo reaction norms*

While we found evidence of additive genetic variance for all monitored traits and under some environmental conditions, we found no indications of heritable variation for reaction norms. As only four males were used in the current study, our power to detect such interactions was limited. Nevertheless, the observed additive genetic variance generally appeared to be context dependent. While certain studies have provided evidence of gene  $\times$  environment interactions on reaction norms, including survival/length (Evans *et al.* 2010b), body mass (Crespel *et al.* 2013), developmental time (Clark *et al.* 2013; Pompini *et al.* 2013) and gene expression (Côté *et al.* 2007) in salmonids, others have not for some of the same traits (larval length: Clark *et al.* 2013; embryonic survival: Pompini *et al.* 2013). These discrepancies may be reflective of differential selective pressures imposed by environmental stressors or of the costs of plasticity varying between species and traits (DeWitt *et al.* 1998).

With respect to gene expression reaction norms, differences may also arise as a consequence of the genes' functions. For example, a study by Landry *et al.* (2006) demonstrated that in yeast, heritable variation in expression reaction norms was biased towards genes with low ties to fitness. This bias was presumably due to the fact that such genes are expected to have fewer constraints on their possible responses to environmental change and therefore have less canalized reaction norms. A wider survey of the transcriptional responses to pathogen stressors would be needed to confirm if the same were true in our whitefish population.

## Conclusions

Treatment of whitefish embryos with two distinct isolates of the opportunistic pathogen *Pseudomonas fluorescens* resulted in decreased expression of MHC class I. The more the MHC gene was transcribed, the later embryos hatched, suggesting a trade-off between expression and an earlier life history transition or the existence of further factors that influence both MHC expression and timing of hatching. Significant heritable variation was found for gene expression, embryo mortality and hatching time under certain experimental conditions, but not for others. However, no evidence of gene by environment interactions on the reaction norms was found for any trait of interest. As heritable variation in reaction norms can play a key role in determining a population's ability to cope with unpredictable environments (Hutchings 2011), its absence could have important implications concerning the whitefish's capacity to adapt to changing ecological conditions.

## Acknowledgements

The authors thank F. Hoffman and P. Tavel for permissions and organizational support, A. Schmid for catching the fish and P. Bize, G. Brazzola, P. Christe, M. Dijkstra, M. dos Santos, L. Fumagalli, F. Glauser, K. Hine, R. Kanitz, M. Lehto, R. Nicolet, M. Pompini, T. Reusch, J.R. van der Meer, C. van Oosterhout and F. Witsenburg for helping in the field and/or discussions. C. Berney and H. Richter helped optimizing the RNA extraction and the qPCR protocols. We also thank L. Bernatchez and five anonymous reviewers for comments on the manuscript. The project was funded by the Swiss National Science Foundation.

## References

- Agrawal A, Conner J, Johnson M, Wallsgrove R (2002) Ecological genetics of an induced plant defense against herbivores: Additive genetic variance and costs of phenotypic plasticity. *Evolution*, **56**, 2206–2213.
- Antoniou AN, Powis SJ (2008) Pathogen evasion strategies for the major histocompatibility complex class I assembly pathway. *Immunology*, **124**, 1–12.
- Arkush KD, Giese AR, Mendonca HL *et al.* (2002) Resistance to three pathogens in the endangered winter-run chinook salmon (*Oncorhynchus tshawytscha*): effects of inbreeding and major histocompatibility complex genotypes. *Canadian Journal of Fisheries and Aquatic Sciences*, **59**, 966–975.
- Aubin-Horth N, Renn SC (2009) Genomic reaction norms: using integrative biology to understand molecular mechanisms of phenotypic plasticity. *Molecular Ecology*, **18**, 3763–3780.
- Austin B, Ausin DA (1999) *Bacterial Fish Pathogens: Disease of Farmed and Wild Fish*. Praxis Publishing Ltd., Chichester.
- Aykanat T, Heath JW, Dixon B, Heath DD (2012) Additive, non-additive and maternal effects of cytokine transcription in response to immunostimulation with *Vibrio* vaccine in

- Chinook salmon (*Oncorhynchus tshawytscha*). *Immunogenetics*, **64**, 691–703.
- Bast E (2001) *Mikrobiologische Methoden: Eine Einführung in Grundlegende Arbeitstechniken*. Spektrum Akademischer Verlag, Berlin.
- Bates D, Maechler M, Bolker B (2011) lme4: Linear mixed-effects models using S4 classes. *R Package Version 0.999375-39*. <http://CRAN.R-project.org/package=lme4>.
- Benedetto A, Squadrone S, Prearo M *et al.* (2011) Evaluation of ABC efflux transporters genes expression in kidney of rainbow trout (*Oncorhynchus mykiss*) fed with melamine and cyanuric acid diets. *Chemosphere*, **84**, 727–730.
- Binz T, Largiadèr C, Müller R, Wedekind C (2001) Sequence diversity of Mhc genes in lake whitefish. *Journal of Fish Biology*, **58**, 359–373.
- Brzuzan P, Jurczyk L, Luczynski MK, Gora M (2005) Relative quantification of CYP1A gene expression in whitefish (*Coregonus lavaretus*) exposed to benzo[a]pyrene. *Environmental Biotechnology*, **1**, 11–15.
- Brzuzan P, Wozny M, Dobosz S *et al.* (2007) Blue sac disease in larval whitefish, *Coregonus lavaretus* (L.): pathological changes in mRNA levels of CYP1A, ER alpha, and p53. *Journal of Fish Diseases*, **30**, 169–173.
- Brzuzan P, Wozny M, Ciesielski S *et al.* (2009) Microcystin-LR induced apoptosis and mRNA expression of p53 and cdkn1a in liver of whitefish (*Coregonus lavaretus* L.). *Toxicol*, **54**, 170–183.
- Bustin SA, Nolan T (2004) Pitfalls of quantitative real-time reverse-transcription polymerase chain reaction. *Journal of Biomolecular Techniques*, **15**, 155–166.
- Cavalieri D, Townsend JP, Hartl DL (2000) Manifold anomalies in gene expression in a vineyard isolate of *Saccharomyces cerevisiae* revealed by DNA microarray analysis. *Proceedings of the National Academy of Sciences of the United States of America*, **97**, 12369–12374.
- Charmantier A, Garant D (2005) Environmental quality and evolutionary potential: lessons from wild populations. *Proceedings of the Royal Society B-Biological Sciences*, **272**, 1415–1425.
- Clark ES, Wedekind C (2011) Additive genetic effects on embryo viability in a whitefish (Salmonidae) influenced by the water mould *Saprolegnia ferax*. *Journal of Bacteriology and Parasitology*, **54**, 1.
- Clark ES, Stelkens RB, Wedekind C (2013) Parental influences on pathogen resistance in brown trout embryos and effects of outcrossing within a river network. *PLoS ONE*, **8**, e57832.
- Côté G, Perry G, Blier P, Bernatchez L (2007) The influence of gene-environment interactions on GHR and IGF-I expression and their association with growth in brook charr, *Salvelinus fontinalis* (Mitchill). *BMC Genetics*, **8**, 87.
- Crespel A, Bernatchez L, Audet C, Garant D (2013) Strain specific genotype-environment interactions and evolutionary potential for body mass in brook charr (*Salvelinus fontinalis*). *G3: Genes Genomes, Genetics*, **3**, 379–386.
- Croisetiere S, Bernatchez L, Bellhumeur P (2010) Temperature and length-dependent modulation of the MH class II beta gene expression in brook charr (*Salvelinus fontinalis*) by a cis-acting minisatellite. *Molecular Immunology*, **47**, 1817–1829.
- Daszak P (2000) Emerging infectious diseases of wildlife - threats to biodiversity and human health. *Science*, **287**, 443–449.
- Debes PV, Normandeau E, Fraser DJ, Bernatchez L, Hutchings JA (2012) Differences in transcription levels among wild, domesticated, and hybrid Atlantic salmon (*Salmo salar*) from two environments. *Molecular Ecology*, **21**, 2574–2587.
- DeWitt TJ, Sih A, Sloan Wilson D (1998) Costs and limits of phenotypic plasticity. *Trends in Ecology & Evolution*, **13**, 77–81.
- Dobson A, Foufopoulos J (2001) Emerging infectious pathogens of wildlife. *Philosophical Transactions of the Royal Society B: Biological Sciences*, **356**, 1001–1012.
- Einum S, Fleming IA (1999) Maternal effects of egg size in brown trout (*Salmo trutta*): norms of reaction to environmental quality. *Proceedings of the Royal Society of London Series B-Biological Sciences*, **266**, 2095–2100.
- Einum S, Fleming I (2000) Selection against late emergence and small offspring in Atlantic salmon (*Salmo salar*). *Evolution*, **54**, 628–639.
- Evans ML, Neff BD, Heath DD (2010a) MHC-mediated local adaptation in reciprocally translocated Chinook salmon. *Conservation Genetics*, **11**, 2333–2342.
- Evans ML, Neff BD, Heath DD (2010b) Quantitative genetic and translocation experiments reveal genotype-by-environment effects on juvenile life-history traits in two populations of Chinook salmon (*Oncorhynchus tshawytscha*). *Journal of Evolutionary Biology*, **23**, 687–698.
- Finlay BB, McFadden G (2006) Anti-immunology: evasion of the host immune system by bacterial and viral pathogens. *Cell*, **124**, 767–782.
- Fischer U, Dijkstra JM, Kollner B *et al.* (2005) The ontogeny of MHC class I expression in rainbow trout (*Oncorhynchus mykiss*). *Fish & Shellfish Immunology*, **18**, 49–60.
- Fisher MA, Oleksiak MF (2007) Convergence and divergence in gene expression among natural populations exposed to pollution. *BMC Genomics*, **8**, 108.
- Frost SDW (1999) The immune system as an inducible defense. In: *The Ecology and Evolution of Inducible Defenses* (eds Tollrian R, Harvell CD), pp. 104–126. Princeton University Press, Princeton.
- Gibson G, Dworkin I (2004) Uncovering cryptic genetic variation. *Nature Reviews Genetics*, **5**, 681–690.
- Haase D, Roth O, Kalbe M *et al.* (2013) Absence of major histocompatibility complex class II mediated immunity in pipefish, *Syngnathus typhle*: evidence from deep transcriptome sequencing. *Biology Letters*, **9**, 20130044.
- Hadfield J (2010) MCMC methods for multi-response generalized linear mixed models: the MCMCglmm R package. *Journal of Statistical Software*, **33**, 1–22.
- Harvell CD, Kim K, Burkholder JM *et al.* (1999) Review: marine ecology - emerging marine diseases - climate links and anthropogenic factors. *Science*, **285**, 1505–1510.
- Heintz RA, Short JW, Rice SD (1999) Sensitivity of fish embryos to weathered crude oil: part II. Increased mortality of pink salmon (*Oncorhynchus gorbuscha*) embryos incubating downstream from weathered Exxon Valdez crude oil. *Environmental Toxicology and Chemistry*, **18**, 494–503.
- Hodgins-Davis A, Townsend JP (2009) Evolving gene expression: from G to E to G x E. *Trends in Ecology & Evolution*, **24**, 649–658.
- Hodgins-Davis A, Adomas AB, Warringer J, Townsend JP (2012) Abundant gene-by-environment interactions in gene expression reaction norms to copper within *Saccharomyces cerevisiae*. *Genome Biology and Evolution*, **4**, 1061–1079.

- Hoffmann A, Merilä J (1999) Heritable variation and evolution under favourable and unfavourable conditions. *Trends in Ecology and Evolution*, **14**, 96–101.
- Houle D (1992) Comparing evolvability and variability of quantitative traits. *Genetics*, **130**, 195–204.
- Hutchings JA (2011) Old wine in new bottles: reaction norms in salmonid fishes. *Heredity*, **106**, 421–437.
- Jacob A, Nusslé S, Britschgi A *et al.* (2007) Male dominance linked to size and age, but not to 'good genes' in brown trout (*Salmo trutta*). *BMC Evolutionary Biology*, **7**, 207.
- Jacob A, Evanno G, von Siebenthal BA, Grossen C, Wedekind C (2010) Effects of different mating scenarios on embryo viability in brown trout. *Molecular Ecology*, **19**, 5296–5307.
- Koho J, Karjalainen J, Viljanen M (1991) Effects of temperature, food density and time of hatching on growth, survival, and feeding of vendace (*Coregonus albula* (L.)) larvae. *Aqua Fennica*, **21**, 63–73.
- Koppang EO, Dannevig BH, Lie O, Ronningen K, Press CM (1999) Expression of Mhc class I and II mRNA in a macrophage-like cell line (SHK-1) derived from Atlantic salmon, *Salmo salar* L., head kidney. *Fish & Shellfish Immunology*, **9**, 473–489.
- Kottelat M, Freyhof J (2007) *Handbook of European Freshwater Fishes*. Publications Kottelat, Cornol, Switzerland, New Jersey.
- Kovacsovics-Bankowski M, Rock KL (1995) A phagosome-to-cytosol pathway for exogenous antigens presented on MHC class-I molecules. *Science*, **267**, 243–246.
- Landry CR, Oh J, Hartl DL, Cavalieri D (2006) Genome-wide scan reveals that genetic variation for transcriptional plasticity in yeast is biased towards multi-copy and dispensable genes. *Gene*, **366**, 343–351.
- Lapaque N, Hutchinson JL, Jones DC *et al.* (2009) Salmonella regulates polyubiquitination and surface expression of MHC class II antigens. *Proceedings of the National Academy of Sciences of the United States of America*, **106**, 14052–14057.
- Larsen PF, Nielsen EE, Williams TD *et al.* (2007) Adaptive differences in gene expression in European flounder (*Platichthys flesus*). *Molecular Ecology*, **16**, 4674–4683.
- Laugen AT, Kruuk LEB, Laurila A *et al.* (2005) Quantitative genetics of larval life-history traits in *Rana temporaria* in different environmental conditions. *Genetical Research*, **86**, 161–170.
- Laurila A, Karttunen S, Merilä J (2002) Adaptive phenotypic plasticity and genetics of larval life histories in two *Rana temporaria* populations. *Evolution*, **56**, 617–627.
- Lenz TL, Becker S (2008) Simple approach to reduce PCR artefact formation leads to reliable genotyping of MHC and other highly polymorphic loci—implications for evolutionary analysis. *Gene*, **427**, 117–123.
- Li Y, Alvarez OA, Gutteling EW *et al.* (2006) Mapping determinants of gene expression plasticity by genetical genomics in *C. elegans*. *Plos Genetics*, **2**, e222.
- Livak KJ, Schmittgen TD (2001) Analysis of relative gene expression data using real-time quantitative PCR and the 2<sup>-</sup>(Delta Delta C(T)) method. *Methods*, **25**, 402–408.
- Lochmiller RL, Deerenberg C (2000) Trade-offs in evolutionary immunology: just what is the cost of immunity? *Oikos*, **88**, 87–98.
- Lynch M, Walsh B (1998) *Genetics and Analysis of Quantitative Traits*. Sinauer Associates Inc, Sunderland, Massachusetts.
- Magnadóttir B (2006) Innate immunity of fish (overview). *Fish Shellfish Immunology*, **20**, 137–151.
- McCairns RJ, Bernatchez L (2010) Adaptive divergence between freshwater and marine sticklebacks: insights into the role of phenotypic plasticity from an integrated analysis of candidate gene expression. *Evolution*, **64**, 1029–1047.
- McGuigan K, Sgro CM (2009) Evolutionary consequences of cryptic genetic variation. *Trends in Ecology and Evolution*, **24**, 305–311.
- Merilä J, Sheldon BC (1999) Genetic architecture of fitness and nonfitness traits: empirical patterns and development of ideas. *Heredity*, **83**, 103–109.
- Merilä J, Sheldon BC, Kruuk LEB (2001) Explaining stasis: microevolutionary studies in natural populations. *Genetica*, **112**, 199–222.
- Merilä J, Soderman F, O'Hara R, Rasanen K, Laurila A (2004) Local adaptation and genetics of acid-stress tolerance in the moor frog, *Rana arvalis*. *Conservation Genetics*, **5**, 513–527.
- Moret Y, Schmid-Hempel P (2000) Survival for immunity: the price of immune system activation for bumblebee workers. *Science*, **290**, 1166–1168.
- Mousseau TA, Roff DA (1987) Natural selection and the heritability of fitness components. *Heredity*, **59**, 181–197.
- Mulero I, Garcia-Ayala A, Meseguer J, Mulero V (2007) Maternal transfer of immunity and ontogeny of autologous immunocompetence of fish: a minireview. *Aquaculture*, **268**, 244–250.
- Nath S, Kales S, Fujiki K, Dixon B (2006) Major histocompatibility class II genes in rainbow trout (*Oncorhynchus mykiss*) exhibit temperature dependent downregulation. *Immunogenetics*, **58**, 443–453.
- Norbury CC, Hewlett LJ, Prescott AR, Shastri N, Watts C (1995) Class I MHC presentation of exogenous soluble antigen via macropinocytosis in bone marrow macrophages. *Immunity*, **3**, 783–791.
- Norris K, Evans MR (2000) Ecological immunology: life history trade-offs and immune defense in birds. *Behavioral Ecology*, **11**, 19–26.
- OECD (1992) *Guideline 203: Fish, Acute Toxicity Test*. Organisation for Economic Co-operation and Development, Paris, France. www.oecd.org.
- Olsvik PA, Lie KK, Jordal AE, Nilssen TO, Hordvik I (2005) Evaluation of potential reference genes in real-time RT-PCR studies of Atlantic salmon. *BMC Molecular Biology*, **6**, 21.
- Pakkasmaa S, Merilä J, O'Hara RB (2003) Genetic and maternal effect influences on viability of common frog tadpoles under different environmental conditions. *Heredity*, **91**, 117–124.
- Pavey SA, Lamaze FC, Garant D, Bernatchez L (2011) Full length MHC IIbeta exon 2 primers for salmonids: a new resource for next generation sequencing. *Conservation Genetic Resources*, **3**, 665–667.
- Perkins JR, Dawes JM, McMahon SB, *et al.* (2012) ReadqPCR and NormqPCR: R packages for the reading, quality checking and normalisation of RT-qPCR quantification cycle (Cq) data. *BMC Genomics*, **13**, 296.
- Phillips RW, Claire EW (2011) Intragravel movement of the reticulate sculpin, *Cottus perplexus* and its potential as a predator on salmonid embryos. *Transactions of the American Fisheries Society*, **95**, 210–212.
- Picard DJ, Schulte PM (2004) Variation in gene expression in response to stress in two populations of *Fundulus heteroclitus*.

- Comparative Biochemistry and Physiology a-Molecular & Integrative Physiology*, **137**, 205–216.
- Pigliucci M (2001) *Phenotypic Plasticity Beyond Nature and Nurture*. The John Hopkins University Press, Baltimore.
- Pitcher TE, Neff BD (2006) MHC class IIB alleles contribute to both additive and nonadditive genetic effects on survival in Chinook salmon. *Molecular Ecology*, **15**, 2357–2365.
- Pompini M, Clark ES, Wedekind C (2013) Pathogen-induced hatching and population-specific life-history response to water-borne cues in brown trout (*Salmo trutta*). *Behavioral Ecology and Sociobiology*, **67**, 649–656.
- R Development Core Team (2011) *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria. <http://www.R-project.org>.
- Relyea RA (2005) The heritability of inducible defenses in tadpoles. *Journal of Evolutionary Biology*, **18**, 856–866.
- Reyes-Becerril M, Lopez-Medina T, Ascencio-Valle F, Esteban MA (2011) Immune response of gilthead seabream (*Sparus aurata*) following experimental infection with *Aeromonas hydrophila*. *Fish & Shellfish Immunology*, **31**, 564–570.
- Rodrigues PNS, Hermsen TT, van Maanen A *et al.* (1998) Expression of *MhcCyca* class I and class II molecules in the early life history of the common carp (*Cyprinus carpio* L.). *Developmental and Comparative Immunology*, **22**, 493–506.
- Runcie DE, Garfield DA, Babbitt CC *et al.* (2012) Genetics of gene expression responses to temperature stress in a sea urchin gene network. *Molecular Ecology*, **21**, 4547–4562.
- Sato A, Figueroa F, Murray BW *et al.* (2000) Nonlinkage of major histocompatibility complex class I and class II loci in bony fishes. *Immunogenetics*, **51**, 108–116.
- Scarpellini M, Franzetti L, Galli A (2004) Development of PCR assay to identify *Pseudomonas fluorescens* and its biotype. *FEMS Microbiology Letters*, **236**, 257–260.
- Schlichting CD, Pigliucci M (1998) *Phenotypic Evolution: A Reaction Norm Perspective*. Sinauer Associates, Sunderland.
- Schmid-Hempel P (2003) Variation in immune defence as a question of evolutionary ecology. *Proceedings of the Royal Society of London Series B-Biological Sciences*, **270**, 357–366.
- Schmittgen TD, Livak KJ (2008) Analyzing real-time PCR data by the comparative C-T method. *Nature Protocols*, **3**, 1101–1108.
- Schreck CB, Contreras-Sanchez W, Fitzpatrick MS (2001) Effects of stress on fish reproduction, gamete quality, and progeny. *Aquaculture*, **197**, 3–24.
- von Siebenthal BA, Jacob A, Wedekind C (2009) Tolerance of whitefish embryos to *Pseudomonas fluorescens* linked to genetic and maternal effects, and reduced by previous exposure. *Fish & Shellfish Immunology*, **26**, 531–535.
- Silver SJ, Warren CE, Doudoroff P (2011) Dissolved oxygen requirements of developing steelhead trout and chinook salmon embryos at different water velocities. *Transactions of the Faraday Society*, **92**, 327–343.
- Skoglund H, Einum S, Forseth T, Barlaup BT (2012) The penalty for arriving late in emerging salmonid juveniles: differences between species correspond to their interspecific competitive ability. *Functional Ecology*, **26**, 104–111.
- Spiers AJ, Buckling A, Rainey PB (2000) The causes of *Pseudomonas* diversity. *Microbiology*, **146**, 2345–2350.
- Star B, Nederbragt AJ, Jentoft S *et al.* (2011) The genome sequence of Atlantic cod reveals a unique immune system. *Nature*, **477**, 207–210.
- Stelkens RB, Pompini M, Wedekind C (2012) Testing for local adaptation in brown trout using reciprocal transplants. *BMC Evolutionary Biology*, **12**, 247.
- Tang J, Bryant MD, Brannon EL (1987) Effect of temperature extremes on the mortality and development rates of coho salmon embryos and alevins. *Progressive Fish-Culturist*, **49**, 167–174.
- Townsend JP, Cavalieri D, Hartl DL (2003) Population genetic variation in genome-wide gene expression. *Molecular Biology and Evolution*, **20**, 955–963.
- Vandesompele J, De Preter K, Pattyn F *et al.* (2002) Accurate normalization of real-time quantitative RT-PCR data by geometric averaging of multiple internal control genes. *Genome Biology*, **3**, 34.
- Warkentin KM (2011) Environmentally cued hatching across taxa: embryos respond to risk and opportunity. *Integrative and Comparative Biology*, **51**, 14–25.
- Wedekind C (2002) Induced hatching to avoid infectious egg disease in whitefish. *Current Biology*, **12**, 69–71.
- Wedekind C, Müller R (2005) Risk-induced early hatching in salmonids. *Ecology*, **86**, 2525–2529.
- Wedekind C, Müller R, Steffen A, Eggler R (2001) A low-cost method of rearing multiple batches of fish. *Aquaculture*, **192**, 31–37.
- Wedekind C, Walker M, Portmann J *et al.* (2004) MHC-linked susceptibility to a bacterial infection, but no MHC-linked cryptic female choice in whitefish. *Journal of Evolutionary Biology*, **17**, 11–18.
- Wedekind C, Evanno G, Urbach D, Jacob A, Müller R (2008) 'Good-genes' and 'compatible-genes' effects in an Alpine whitefish and the information content of breeding tubercles over the course of the spawning season. *Genetica*, **134**, 21–30.
- Wedekind C, Gessner MO, Vazquez F, Maerki M, Steiner D (2010) Elevated resource availability sufficient to turn opportunistic into virulent fish pathogens. *Ecology*, **91**, 1251–1256.
- Wilson AJ, Pemberton JM, Pilkington JG *et al.* (2006) Environmental coupling of selection and heritability limits evolution. *PLoS Biology*, **4**, e216.
- Windsor DA (1998) Most of the species on Earth are parasites. *International Journal for Parasitology*, **28**, 1939–1941.
- Yewdell JW, Norbury CC, Bennink JR (1999) Mechanisms of exogenous antigen presentation by MHC class I molecules *in vitro* and *in vivo*: implications for generating CD8(+) T cell responses to infectious agents, tumors, transplants, and vaccines. *Advances in Immunology*, **73**, 1–77.
- Zapata A, Diez B, Cejalvo T, Gutierrez-de Frias C, Cortes A (2006) Ontogeny of the immune system of fish. *Fish & Shellfish Immunology*, **20**, 126–136.
- Zhang WW, Hu YH, Wang HL, Sun L (2009) Identification and characterization of a virulence-associated protease from a pathogenic *Pseudomonas fluorescens* strain. *Veterinary Microbiology*, **139**, 183–188.

---

E.C., L.W., and C.W. designed the experiment. E.C. and L.W. executed the experiment, and E.C., L.W., and C.W. analysed the data and wrote the manuscript.

---

### Data accessibility

Data package with one file including hatching success and time until hatching for each monitored embryo and one file including  $2^{-\Delta CT}$  values for each gene and embryo: DRYAD entry doi:10.5061/dryad.5sp75.

### Supporting information

Additional supporting information may be found in the online version of this article.

**Table S1** Mean MHC class I expression levels for each full-sib family and associated variances.

**Table S2** Posterior mean estimates and 95% highest posterior density confidence intervals for the effects of treatment on

MHC I expression, embryonic mortality, and hatching time, as estimated with MCMCglmm.

**Table S3** Posterior modes of variance estimates for MHC class I expression, embryonic mortality, and hatching time in each treatment (control, PF1, and PF2), as estimated with MCMCglmm.

**Table S4** Comparison of deviance information criterion (DIC) of the reference model (in bold) to models incorporating interaction effects, as estimated with MCMCglmm.

**Fig. S1** Representative agarose gels showing amplification of MHC class I (A-B) and class II (C-D) from the cDNA of four adult whitefish spleens (A, C) and four whole embryos (B, D).

**Fig. S2** Treatment effects on hatching time (means  $\pm$  SE) in a (A) simulated data set and a (B) data set including only full-sib families with no embryonic mortality.